



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

In order to determine the individual variations of plants taken at the same time of year and treated in exactly the same way, the following species were studied with appended results:

April, 5 plants *Pelargonium zonale*, .620, .645, .767, .782, 1.035 atmospheres.
June, 5 plants *Fuchsia speciosa*, 1.406, 1.428, 1.467, 1.500, 1.605 “

While an average of the pressures given by the above plants may seem of no great value, nevertheless there is a certain advantage in knowing this mean, which may be taken as a general expression of the root pressure of common greenhouse plants. The mean pressure of the 22 plants studied is approximately 0.9 atmosphere, or in round numbers 13^{lb} for the square inch.

Turning from pressure to quantity of exudation, the above table shows that the plants giving the greatest quantity of exudation are *Fuchsia speciosa* and *Begonia coccinea*. Different plants of *Fuchsia speciosa* vary greatly in the quantity given off, and this seems to be correlated with the formation of new shoots—the longer this is delayed the greater the flow. This is true in less degree of most of the plants which form new shoots. In others—*Impatiens Holstii*, *Lycopersicum esculentum*, *Pelargonium peltatum*, and *Tropaeolum majus*—the stems decay and the roots soon die.

Some observations were made on periodicity of exudation and of exudation pressure. These facts will be published in a separate note later.—
SOPHIA ECKERSON, *Northampton, Mass.*

THE CONDITION OF CERTAIN WINTER BUDS

During 1905-1906 an effort was made to ascertain the seasonal stages in the microspore development of certain woody plants, with the following results:

	Mother cell stage	Dividing nuclei	Mature spores
<i>Populus deltoides</i>	October 23
<i>Fraxinus americana</i>	Oct. 23—Feb. 15
<i>Celtis occidentalis</i>	October 24	April 12	April 15
<i>Carpinus caroliniana</i>	October 24	April 12	April 15
<i>Cornus florida</i>	August 20	August 22	August 24
<i>Cercis canadensis</i>	October 31	April 17	April 20

WILLIAM MOORE and MARY E. BEHNEY, *Ursinus College, Pa.*